



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

AVERAGE AND PROBABILITY.

85. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

Two points are taken at random in a circle and a chord drawn through them; a point is then taken at random in each segment. Find the average area of the quadrilateral formed by joining the four points.

86. Proposed by L. C. WALKER, Assistant Professor of Mathematics, Leland Stanford, Jr., University, Palo Alto, Cal.

Two points are taken at random in a circular annulus formed by two concentric circles. Find the chance that the straight line joining the points will not cut the inner variable circle.

87. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

Find the mean distance of a random point in a sphere from a point, (1) within, (2) without the sphere.

** Solutions of these problems should be sent to B. F. Finkel not later than January 10.

EDITORIALS.

Prof. James Pierpont, of Yale University, was elected a member of the French Mathematical Society.

Dr. Artemas Martin, of the United States Coast and Geodetic Survey, at Washington, D. C., has recently been elected a member of the Deutsche Mathematiker-Vereinigung.

At the recent meeting of the Board of Trustees of the Indiana University, Dr. John B. Faught, Instructor in Mathematics, was given the title of Assistant Professor of Mathematics.

Prof. J. H. Tanner, of Cornell University, is writing an elementary algebra for the "Cornell Series," and Prof. James McMahon is writing an elementary geometry for the same series.

We are pleased to state that Dr. Halsted is receiving many congratulatory letters from readers of the **MONTHLY** who are reading with interest and profit, his contributions to the **MONTHLY** in Non-Euclidean Geometry.

Through the kindness of Dr. Alexander Macfarlane, we have received for publication a paper on "The Theory of Mathematical Inference," by Prof. G. J. Stokes, of Queen's College, Cork, Ireland. This paper will appear in the January number, Vol. VII, of the **MONTHLY**.

A number of our readers have asked us to send them the July issue. In answer we will say, take the cover off the June number and you will then have a June-July number. Through error on the part of the printers the statement of a double number was omitted on the cover.